

# Particulate Matter Emissions from Household Cooking Methods and Associated Respiratory Health Risks: A Systematic Review

Ruthvik Yaparla

Round Rock, 201 Deep Wood Dr, Round Rock, TX, 78681, USA

## ABSTRACT

Indoor air quality remains largely unregulated although people spend an average of 21.6 hours indoors daily, compared to only 2.4 hours outdoors. While ambient air pollution has been extensively studied, there is a significant gap in research on household and indoor air pollution, particularly regarding emissions from everyday cooking practices. This study explores commonly used cooking methods, their associated particulate matter (PM) emissions, their potential impacts on respiratory health, and gaps in current literature. A total of 416 unique studies were screened, with 354 excluded based on title and abstract. Sixty-two reports were sought for retrieval, and ultimately 13 studies were included in the review. The results revealed that pan-frying generated a mean PM1 concentration of 139.55  $\mu\text{g}/\text{m}^3$ , while toasting produced 58.8  $\mu\text{g}/\text{m}^3$ . Deep-frying yielded the highest mean PM2.5 concentration at 841  $\mu\text{g}/\text{m}^3$ , followed by stewing, stir-frying, roasting, pan-frying, boiling, toasting, and steaming, in descending order. Similarly, deep-frying produced the highest PM10 levels (1192  $\mu\text{g}/\text{m}^3$ ) with roasting, pan-frying and toasting also contributing substantially. Interestingly, when comparing frying and non-frying methods, no statistically significant differences in emission levels were found. However, this analysis is limited by the generalization of data across regions and cooking cultures, as variations in oils and fats may influence emission profiles. Despite these limitations, this study establishes a foundational understanding of PM emissions by cooking method and their potential links to respiratory and cardiovascular health outcomes, lung function, and carcinogen exposure. Further research is warranted to evaluate indoor air pollution associated with specific appliances and cooking methods, particularly emerging technologies such as air fryers.

**Keywords:** Particulate Matter; Pollution; Respiratory; Lungs; Disease; Cooking; Indoors

## INTRODUCTION

Ambient (outdoor) air pollution was first identified as a global health concern decades ago and governments have been implementing successful measures to mitigate these effects since then. In the next few years, ambient air pollution is predicted to decrease substantially. But what about indoor air pollution? The Global Burden of Disease study concluded that in 2021, outdoor and indoor

---

**Corresponding author:** Ruthvik Yaparla, E-mail: [ruthvikyaparla24@gmail.com](mailto:ruthvikyaparla24@gmail.com)

**Copyright:** © 2025 Ruthvik Yaparla. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Received** June 3, 2025; **Accepted** July 3, 2025

<https://doi.org/10.70251/HYJR2348.3419>

air pollution contributed to 8 million deaths worldwide (1). Poor indoor air quality (IAQ) is a significant factor in deaths from air pollution and its contribution is often overlooked. An article in the journal *Nature* stated, “Researchers and policymakers are only now waking up to the effects of dirty indoor air. As ever, low-income and marginalized communities are most exposed” (2).

According to a survey conducted by the United States Environmental Protection Agency (EPA), humans spend 90% of their time indoors and only 10% outdoors (3). Furthermore, elderly people, a demographic that tends to be more vulnerable to respiratory complications, spend an average of 92% of their day indoors (4). Yet, significantly less research has been undertaken regarding IAQ compared to ambient air quality. Previous research has investigated the effects of a culmination of various indoor air pollutants, such as incense burning, vacuuming, and candles. However, cooking emissions are rarely ever the sole focus of a study, even though people spend, on average, a greater amount of their day cooking than all these other activities. According to the US Bureau of Labor Statistics, Americans spend an average of an hour preparing food every day using cooking appliances that are known to release harmful chemicals (5).

Significant research has demonstrated the effects of Volatile Organic Compounds (VOCs) without a strong focus on particulate matter (PM), even though PM is typically more harmful. Outdoor PM is considered a Group 1 Carcinogen by the International Agency for Research on Cancer (IARC) (6). On the other hand, indoor PM isn’t considered a carcinogen. This is a concern because even though outdoor air pollution contains harmful chemicals and drives climate change, poor IAQ exposes humans to greater concentrations of PM, which may elevate the risk for health complications. Three types of PM will be investigated in this review: PM<sub>10</sub>, describing particles with a diameter of 10 µm or smaller, PM<sub>2.5</sub>, describing particles with a diameter of 2.5 µm or smaller, and PM<sub>1</sub>, describing the smallest size with particles of diameter 1 µm or smaller.

There is significant understanding regarding types of PM prevalent in the outdoor environment, however, the gaps in understanding the cooking methods that contribute most to PM emissions in an indoor environment shouldn’t be ignored. Resolving these gaps is important because confined spaces, such as kitchens, can cause PM to accumulate in great concentrations and cause respiratory and cardiovascular disease. This systematic review and aggregated means analysis will investigate the overlooked role of cooking emissions on respiratory risk exposure

and can serve as a guideline for future research on the subject matter.

## **METHODS**

### **Database and Search Strategy**

Online searches were completed using the PubMed database. This systematic review followed guidelines set by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement (7). Published items relevant to the research question, “How do Cooking Methods’ Release of Particulate Matter Contribute to Respiratory Risk Exposure?”, were assessed. A boolean search algorithm was implemented using the terms “carcinogen,” “air quality,” “particulate matter,” and “respiratory exposure,” in conjunction with the names of specific common cooking methods. Boolean expressions “AND” and “OR” were used alongside these search terms as needed. A total of 16 searches were conducted. Finally, the “10 years” filter was applied to all searches to retrieve up-to-date studies. The search was performed from February 2024 to June 2024.

### **Inclusion Criteria**

The “10 years” filter was applied to all PubMed searches to retrieve studies that have been published within the last 10 years. This is because studies from over 10 years ago often lacked access to air filtration and ventilation that are common today. This, in turn, would cause the PM emission concentrations from those studies to be abnormally high. Published items such as review articles, abstracts, and articles not written in English were excluded.

### **Selection of Studies**

Initial records were screened for eligibility based on title. After duplicate records were removed, articles were excluded based on abstract. Finally, the remaining articles were full-text assessed for the relevancy of data points. Reports containing data points such as PM emission concentrations from specific cooking methods were then extracted and compiled (Figure 1) (Tables 1 and 2).

### **Statistical Analysis**

The mean emission concentrations of PM<sub>1</sub>, PM<sub>2.5</sub>, and PM<sub>10</sub> were extracted from all eligible studies. Eligible studies must have recorded PM emission concentrations from specific cooking methods. Furthermore, the reported emissions must not have been emission rates or emission factors. The final cooking methods selected for statistical

analysis were deep-frying, stir-frying, boiling, pan-frying, toasting, roasting, stewing, and steaming. All the PM emission data from eligible studies were organized into groups based on the cooking method and were further separated based on the size of PM reported. The majority of the reported concentrations were in units of  $\mu\text{g}/\text{m}^3$  (micrograms per cubic meter), but some studies reported emission concentrations in units of  $\text{mg}/\text{m}^3$  (milligrams per cubic meter). The data from these studies were converted to units of  $\mu\text{g}/\text{m}^3$ . Then, the mean emission concentrations for PM1, PM2.5, and PM10 were calculated for each cooking method.

After compiling the mean PM emission concentrations of the three sizes for each cooking method, an independent (unpaired), two-tailed Student's t-test was conducted

between two groups for PM2.5 and PM10. The two groups were cooking methods that involve frying (deep-frying, stir-frying, pan-frying) and cooking methods that don't involve frying (boiling, stewing, steaming, roasting, toasting). This test was conducted to see if there was a statistically significant difference in the emission concentrations of PM2.5 and PM10 between the two groups.

## RESULTS

### Mean PM Emission Concentrations

Only one included study reported PM1 emission concentrations for specific cooking methods. Data was extracted from Soppa et al. 2014, which is the only available




Figure 1. Flow chart of the literature search.

Table 1. Summary of included studies [first six studies of the thirteen studies included in the review]

Reference and year	Location/Setting	Number of subjects	Mean PM1 emission concentrations	Mean PM2.5 emission concentrations	Mean PM10 emission concentrations	Cooking method(s)	Health Complications/Respiratory Symptoms
Siponen et al, 2019	Suburb in Kuopio, Finland	37 elderly residents	Not stated	pan frying: 4.4 µg/m <sup>3</sup>	Not stated	Pan Frying	Not stated
			Toasting Bread Level 1*: 37.7 ± 7.0 µg/m <sup>3</sup> ; Level 2*: 79.9 ± 16.1 µg/m <sup>3</sup>		Toasting Bread Level 1: 125.6 ± 87.1 µg/m <sup>3</sup> ; Level 2: 84.6 ± 18.6 µg/m <sup>3</sup>		
			toasting bread Level 1: 62.6 ± 27.7 µg/m <sup>3</sup> ; Level 2: 81.6 ± 16.6 µg/m <sup>3</sup>				
Soppa et al, 2014	Not stated	55 healthy individuals( 28 men + 27 women)	Pan Frying Level 1: 71.3 ± 28.2 µg/m <sup>3</sup> ; Level 2: 207.8 ± 62.4 µg/m <sup>3</sup>	pan frying Level 1: 84.4 ± 37.3 µg/m <sup>3</sup> ; Level 2: 235.2 ± 81.4 µg/m <sup>3</sup>	Pan Frying Level 1: 100.0 ± 51.9 µg/m <sup>3</sup> ; Level 2: 296.9 ± 133.9 µg/m <sup>3</sup>	Toasting Pan Frying	"Our study indicates a possible association of short-term exposure to fine and ultrafine particles emitted from common indoor sources with small decreases in lung function in healthy adults."
			Chinese stall: 201.8 ± 140.5 µg/m <sup>3</sup>				
See et al, 2006	National University of Singapore (NUS) Kent Ridge campus	NA	Not stated	Malay stall: 245.3 ± 77.1 µg/m <sup>3</sup> Indian stall: 186.9 ± 43.6 µg/m <sup>3</sup>	Not stated	Deep Frying Stir Frying Simmering	PM2.5 levels measured at the 3 food stalls far exceeded the National Ambient Air Quality Standard (NAAQS) at 65 µg/m <sup>3</sup> for the 24hr standard.
Sjaastad et al, 2010	Model kitchen in conditions similar to those in a Western European restaurant kitchen	NA	Not stated	Not stated	Not stated	Pan Frying on electric stove Pan Frying on gas stove	Not stated
Sofuoğlu et al, 2015	A Turkish university canteen that serves the school of architecture	NA	Not stated	deep frying: 108 ± 44 µg/m <sup>3</sup> deep-frying: 1720 µg/m <sup>3</sup> stir-frying: 629.33 µg/m <sup>3</sup> stewing: 573 µg/m <sup>3</sup> quick-frying: 201 µg/m <sup>3</sup> boiling: 125.5 µg/m <sup>3</sup> steaming: 49.5 µg/m <sup>3</sup>	1192 µg/m <sup>3</sup>	Deep Frying Deep Frying Stir Frying Stewing Quick Frying Boiling Steaming	"PM10 concentrations associated with frying are much higher than the indoor air standard level of 8-h average 180 µg/m <sup>3</sup> in Hong Kong but lower than the occupational standard of 8-h average 10,000 µg/m <sup>3</sup> set by ACGIH, but lower than the occupational standard of 8-h average 10,000 µg/m <sup>3</sup> set by ACGIH."
Lu et al, 2019	Nankai District in Tianjin, China	NA	Not stated	Not stated	Not stated	Not stated	Not stated

Table 2. Summary of included studies [the remaining 7 studies included in the review]

Reference and year	Location/Setting	Number of subjects	Mean PM1 emission concentrations	Mean PM2.5 emission concentrations	Mean PM10 emission concentrations	Cooking method(s)	Health Complications/Respiratory Symptoms
Lee et al, 2020	Republic of Korea(South Korea)	NA	Not stated	grilling: 754 mg-PM <sub>10</sub> -kg-meat deep-frying: 695.0 µg/m <sup>3</sup> pan-frying: 646.3 µg/m <sup>3</sup> stir-frying: 487.9 µg/m <sup>3</sup> boiling: 138.5 µg/m <sup>3</sup> steaming: 31.3 µg/m <sup>3</sup>	Not stated	Grilling Deep Frying Pan Frying Stir Frying Boiling Steaming	Not stated
Sharma et al, 2020	Residential apartment in the western region of Singapore	NA	Not stated		Not stated		Not stated
Kong et al, 2020	Namdong District, South Korea	NA	Not stated	roasting: 96.21 ± 65.72 µg/m <sup>3</sup>	127.01 ± 119.26 µg/m <sup>3</sup>	Roasting	Not stated
Lee et al, 2022	Schools in Republic of Korea(South Korea)	NA	oily cooking: 13.30 µg/m <sup>3</sup> little oily cooking: 9.80 µg/m <sup>3</sup>	oily cooking: 15.30 µg/m <sup>3</sup> little oily cooking: 11.20 µg/m <sup>3</sup>	oily cooking: 17.0 µg/m <sup>3</sup> little oily cooking: 12.90 µg/m <sup>3</sup>	Oily Cooking(deep frying, stir frying, roasting) Little Oily Cooking(boiling, steaming)	Not stated
Le et al, 2022	Kitchen of commercial Chinese restaurant	NA	Not stated	deep frying: 0.69 ± 0.11 mg/min grilling: 1.58 ± 0.25 mg/min	Not stated	Deep Frying Grilling	"These compounds included known carcinogens such as benzene, formaldehyde, acrolein, and acetaldehyde at concentrations exceeding the RIC limits set by the US EPA and ATSDR."
Padhi et al, 2022	Indo-Gangetic Plain(Uttar Pradesh)	NA	Not stated	boiling: 7.0 ± 2.7 g/kg	Not stated	Boiling	Not stated
Lyu et al, 2022	6 types of commercial kitchens in Shanghai	Not stated	Not stated	teppanyaki kitchen: 850.4 ± 533.4 µg/m <sup>3</sup> barbecue kitchen: 146.6 ± 59.8 µg/m <sup>3</sup> chinese kitchen: 679.1 ± 1922.5 µg/m <sup>3</sup> western kitchen: 272.2 ± 250.1 µg/m <sup>3</sup> fried chicken kitchen: 257.2 ± 44.5 µg/m <sup>3</sup> hotpot cooking area: 110.8 ± 59.8 µg/m <sup>3</sup>	Not stated	Teppanyaki Kitchen Barbecue Kitchen Chinese Kitchen Western Kitchen Fried Chicken Kitchen Hotpot Cooking Area	Chinese kitchens (CK) have the highest deposition rates across all regions of the respiratory system including the upper airways, tracheobronchial, and alveolar regions. Furthermore, the alveolar region generally has the highest deposition rates compared to all other regions, indicating that PM <sub>2.5</sub> penetrates deep into the lungs.

study that reported the PM1 emissions from pan-frying sausages and toasting (8). This study reported emission concentrations for two different exposure levels for both cooking methods. The mean PM1 emission concentrations were calculated to be  $139.55\mu\text{g}/\text{m}^3$  and  $58.8\mu\text{g}/\text{m}^3$  for pan-frying and toasting, respectively (Figure 2).

All eight selected cooking methods had enough data to be considered for statistical analysis of PM2.5 emissions. Deep-frying produced the highest observed mean PM2.5 concentration at  $841\mu\text{g}/\text{m}^3$ , followed by stewing ( $573\mu\text{g}/\text{m}^3$ ), stir-frying ( $558.615\mu\text{g}/\text{m}^3$ ), roasting ( $461.375\mu\text{g}/\text{m}^3$ ), pan-frying ( $234.26\mu\text{g}/\text{m}^3$ ), boiling ( $132\mu\text{g}/\text{m}^3$ ), toasting ( $72.1\mu\text{g}/\text{m}^3$ ), and steaming ( $40.4\mu\text{g}/\text{m}^3$ ). (Figure 2). Finally, for PM10, the mean emission concentrations in order of highest observed value to lowest observed value was as follows: deep frying with a mean concentration of  $1192\mu\text{g}/\text{m}^3$ , roasting with a mean concentration of  $736.99\mu\text{g}/\text{m}^3$ , pan-frying with a mean concentration of  $198.45\mu\text{g}/\text{m}^3$ , and toasting with a mean concentration of  $105.1\mu\text{g}/\text{m}^3$  (Figure 2).

### Comparison of PM2.5 and PM10 Emissions From Frying Versus Non-frying Cooking Methods


The null hypothesis for the t-test analysis of PM2.5 and PM10 emissions between frying cooking methods (deep-frying, stir-frying, pan-frying) and non-frying cooking methods (boiling, stewing, steaming, roasting, toasting) was as follows: “There is no statistically significant difference between mean emissions of cooking methods that involve frying and cooking methods that don’t involve frying.” For PM2.5 emissions between the two study groups,  $p > 0.05$ , indicating a lack of a

statistically significant difference between the amount of PM2.5 emissions from frying and non-frying methods. Only four cooking methods were involved in the t-test for PM10 emissions. Two methods involved frying (deep-frying and pan-frying) and two methods didn’t involve frying (roasting and toasting). The difference in mean PM10 emissions between the two study groups had a  $p > 0.05$ , indicating the absence of a statistically significant difference between the amount of PM10 emissions from frying and non-frying methods. A t-test analysis was not conducted for PM1 emissions between frying and non-frying methods due to a lack of data from eligible studies.

## DISCUSSION

The U.S. Environmental Protection Agency (EPA) has two primary standards for fine particulate matter (PM2.5): an annual average of 9.0 micrograms per cubic meter and a 24-hour average of 35 micrograms per cubic meter (9). The EPA strengthened the primary annual PM2.5 standard from 12.0 to 9.0 micrograms per cubic meter on February 7, 2024. These standards state that, on average, the concentration of PM2.5 should not exceed  $9.0\mu\text{g}/\text{m}^3$  when measured across the span of a year. In addition, on any given day, the concentration of PM2.5 shouldn’t exceed  $35\mu\text{g}/\text{m}^3$ . However, the data from this aggregated means analysis shows that PM2.5 emissions from every selected cooking method significantly exceeded these standards (Figure 2). The EPA has a similar 24-hour standard for PM10: an area meets the standard if it does not exceed a concentration of  $150\mu\text{g}/\text{m}^3$  more than once per year on average over three years (10). Once again, the PM10 emissions from every selected study in this aggregated means analysis exceeded this standard significantly (Figure 2). This raises concern given that there are no current regulations of IAQ in households that use these cooking methods daily.

Unlike PM2.5 and PM10, PM1 is unregulated, and monitoring technology for PM1 is lacking (11). The specific harm that PM1 causes, in contrast to other particulate pollutants and ultrafine particles, is still under investigation (11). When calculating the increased risk of cardiovascular disease from three types of PM, Yin and colleagues found a 0.29% increase for every  $10\mu\text{g}/\text{m}^3$  increase in PM1, which was significantly greater than the increases in cardiovascular disease from PM2.5 and PM10 exposure (12). This study supports the idea that PM1 may be a higher risk factor for disease due to its small size, allowing it to penetrate deeper into vessels than other sizes of PM. This warrants more research to better



**Figure 2.** Mean PM1, PM2.5, and PM10 emissions from different cooking methods as retrieved from the included studies.

understand the relationship between PM1 emissions and respiratory and cardiovascular health.

Some studies found that PM exposure from specific cooking methods was associated with detrimental health effects. For example, Soppa et al. measured forced expiratory volume (FEV), which is the amount of air a person can forcefully exhale after a deep breath, to determine the lung function of healthy adults before and after exposure to PM from toasting bread and pan-frying (8). They concluded that a 10  $\mu\text{g}/\text{m}^3$  increase in PM10, PM2.5, and PM1 emitted from pan-frying sausages was associated with decreases in FEV. On the other hand, they didn't identify any associations between toasting bread and changes in lung function. Le et al. found that Benzene, Formaldehyde, Acrolein, and Acetaldehyde were among the compounds found within emissions from deep-frying and grilling (13). Furthermore, concentrations of Benzene, Acrolein, and Formaldehyde significantly exceeded the Reference Concentrations (RfCs) limits set by the U.S. EPA, and concentrations of Formaldehyde exceeded Minimal Risk Levels (MRLs) set by the Agency for Toxic Substances and Disease Registry (ATSDR). Benzene, Formaldehyde, and Acetaldehyde are all classified as Group 1 Carcinogens by the International Agency for Research on Cancer, while Acrolein is classified as a Group 2 Carcinogen (14). This supported the notion that deep-frying may lead to the highest elevated risk level for health complications amongst all cooking methods because not only does it release the highest observed concentrations of PM, but it also contains significant amounts of carcinogenic compounds.

This review considered each cooking method as either frying (Group 1) or non-frying (Group 2) to conduct a t-test investigating whether there is a significant difference in the amount of emissions between the two groups for PM2.5 and PM10 (PM1 was excluded due to lack of available data). The results showed a lack of statistical evidence to support that one group emits a greater concentration of particles than the other. However, this categorization may oversimplify complex PM emission profiles. For instance, roasting often involves high temperatures and frying fats, which are characteristic of frying cooking methods, leading to greater concentrations of particle emissions compared to other non-frying methods, making it more difficult to conclude a statistically significant difference in emission concentrations between frying and non-frying methods. The uncertainty in data showcases the caution required when approaching such data because variables such as temperature and type of frying fat are inconsistent among different studies.

There is an urgent need for further research regarding PM emissions from air fryers and their effect on respiratory health. Air fryers are relatively new and quickly becoming one of the most popular cooking appliances. However, there is a significant lack of research on how air fryers may elevate the risk level of respiratory disease compared to other cooking methods. An experiment by Wang et al. 2023 found that PM10 emissions from the air frying of chicken wings and breast were higher than pan cooking by a factor of 2.1 and 5.4, respectively (15). This data may suggest that air frying causes a significant elevation in risk level because this review found that, amongst selected studies, pan-frying was the most polluting method in terms of PM1 emissions, the fifth most polluting for PM2.5 emissions, and the third most polluting for PM10 emissions. The conclusions from Wang et al. show the possibility for air frying being one of the most polluting cooking methods.

## LIMITATIONS

Acknowledging that only 13 studies were included in this review to obtain particulate matter emission data, the limitations must be clearly outlined. It is important to avoid drawing significant relationships in the data due to the small sample size of studies. However, this issue directly results from the lack of available literature regarding particulate matter emissions from common cooking appliances. Specifically, only 1 study showcasing PM1 concentrations was deemed eligible for use in this review, even after screening hundreds of articles regarding particulate matter emissions. Using this single study to conclude the severity of the effects that PM1 may cause is dangerous.

In addition, sample heterogeneity may be prevalent in the form of differences in cooking practices between separate regions and cultures. Different cultures implement different frying fats and oils, which could have led to variability in the collected data. For instance, Sjaastad et al. found that pan-frying on an electric stove with margarine as the frying fat had a total particle emission of 1.8mg/m<sup>3</sup>, while cooking with soybean oil as the frying fat had a total particle emission of 1.6mg/m<sup>3</sup> (16). In the same experiment, they reported a total particle emission of 5.5mg/m<sup>3</sup> for pan-frying on a gas stove with margarine as the frying fat, whereas cooking with soybean oil had an emission report of 7.2mg/m<sup>3</sup>. So, study methods that involve the use of a specific oil or frying fat may have different emission data from studies that used other oils and frying fats.

Another important gap that serves as a limitation to this review is that the assays used to measure emissions were inconsistent between selected studies. Given the high variations in PM concentrations reported, further research must be undertaken regarding the quality of methods implemented for measuring particle emissions in indoor settings. For instance, different studies located in different regions used varying equipment and methods to measure emission concentrations, which could have resulted in dramatic variability in the collected data. This makes it difficult to generalize the results of this systematic review across different regions and cooking cultures.

## CONCLUSION

This review synthesized data from 13 studies and found that PM emissions from common cooking methods often exceeded the standards set by the EPA. Furthermore, associations between cardiovascular disease, detriments to lung function, and the presence of significant amounts of carcinogenic compounds were identified from the emissions of several cooking methods. The impact of such emissions from common cooking methods is often overshadowed by the types of cooking fuel used. Future research must investigate the health impacts of specific cooking methods as opposed to solely the type of fuel. Furthermore, in future research, it's important to keep specific variables constant, such as the measuring methods and equipment, as well as the types of oil and frying fat applied when measuring emissions.

## ACKNOWLEDGEMENTS

Agastya Vaidya, Albert Einstein College of Medicine.

## DECLARATION OF CONFLICT OF INTERESTS

The author declares that there are no conflicts of interest regarding the publication of this article.

## REFERENCES

1. IHME, Global Burden of Disease (2024) – with minor processing by Our World in Data. “Alcohol use disorders” [dataset]. IHME, Global Burden of Disease, “Global Burden of Disease - Deaths and DALYs” (accessed on 2024-06-28).
2. “Indoor Air Pollution Kills and Science Needs to Step Up.” Nature News, Nature Publishing Group, 8 Feb. 2023, www.nature.com/articles/d41586-023-00338-0#:~:text=Researchers%20and%20policymakers%20are%20only,marginalized%20communities%20are%20most%20exposed.&text=The%20image%20of%20air%20pollution,can%20be%20a%20misleading%20picture (accessed on 2024-07-05).
3. Klepeis NE, Nelson WC, Ott WR, Robinson JP, *et al.* The National Human Activity Pattern Survey (NHAPS): a resource for assessing exposure to environmental pollutants. *J Expo Anal Environ Epidemiol.* 2001 May-Jun; 11 (3): 231-52. doi: 10.1038/sj.jea.7500165. PMID: 11477521.
4. Siponen T, Yli-Tuomi T, Tiittanen P, Taimisto P, *et al.* Wood stove use and other determinants of personal and indoor exposures to particulate air pollution and ozone among elderly persons in a Northern Suburb. *Indoor Air.* 2019 May; 29 (3): 413-422. doi: 10.1111/ina.12538. Epub 2019 Feb 20. PMID: 30790356; PMCID: PMC6850052.
5. “Exploring time spent on cooking, reading, and other activities for National Hobby month.” U.S. Bureau of Labor Statistics, <https://www.bls.gov/opub/ted/2024/exploring-time-spent-on-cooking-reading-and-other-activities-for-national-hobby-month.htm> (accessed on 2024-07-03).
6. “Traffic-Related Outdoor Air Pollution - NCI.” Dceg.cancer.gov, 15 July 2015, [dceg.cancer.gov/research/what-we-study/traffic-related-outdoor-air-pollution#:~:text=The%20International%20Agency%20for%20Research](https://dceg.cancer.gov/research/what-we-study/traffic-related-outdoor-air-pollution#:~:text=The%20International%20Agency%20for%20Research) (accessed on 2024-07-03).
7. PRISMA. “Welcome to the NEW Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) Website.” [www.prisma-statement.org](http://www.prisma-statement.org), 2020, [www.prisma-statement.org/](http://www.prisma-statement.org/) (accessed on 2024-06-28).
8. Soppa VJ, Schins RP, Hennig F, Hellack B, *et al.* Respiratory effects of fine and ultrafine particles from indoor sources--a randomized sham-controlled exposure study of healthy volunteers. *Int J Environ Res Public Health.* 2014 Jul 4; 11 (7): 6871-89. doi: 10.3390/ijerph110706871. PMID: 25000149; PMCID: PMC4113851.
9. US EPA. “National Ambient Air Quality Standards (NAAQS) for PM.” US EPA, 13 Apr. 2020, [www.epa.gov/pm-pollution/national-ambient-air-quality-standards-naaqs-pm](http://www.epa.gov/pm-pollution/national-ambient-air-quality-standards-naaqs-pm) (accessed on 2024-07-08).
10. The National Ambient Air Quality Standards for Particle Pollution, “Revised Air Quality Standards For Particle Pollution And Updates To The Air Quality Index” (AQI), 2012. (Accessed on 2024-07-11).
11. “IQAir | First in Air Quality.” [www.iqair.com](http://www.iqair.com), [www.iqair.com/us/newsroom/pm1](http://www.iqair.com/us/newsroom/pm1) (accessed on 2024-07-11).
12. Yin Peng, *et al.* Higher Risk of Cardiovascular Disease Associated with Smaller Size-Fractioned Particulate Matter. *Environmental Science & Technology Letters.* 2020; 7 (2): 95–101. <https://doi.org/10.1021/acs.estlett.9b00735>. Accessed 28 Apr. 2021.
13. Le YT, Youn JS, Cho H, Jeon K, *et al.*  $\alpha$ -Fe<sub>2</sub>O<sub>3</sub> nanoparticles and hazardous air pollutants release during cooking using cast iron wok in a commercial Chinese restaurant.

- Environ Pollut. 2022 Aug 15; 307: 119578. doi: 10.1016/j.envpol.2022.119578. Epub 2022 Jun 7. PMID: 35688388.
14. "Known and Probable Human Carcinogens." Www.cancer.org, www.cancer.org/cancer/risk-prevention/understanding-cancer-risk/known-and-probable-human-carcinogens.html (accessed on 2024-07-11).
  15. Wang X, Chan AWH. Particulate Matter and Volatile Organic Compound Emissions Generated from a Domestic Air Fryer. *Environ Sci Technol*. 2023 Nov 14; 57 (45): 17384-17392. doi: 10.1021/acs.est.3c04639. Epub 2023 Nov 6. PMID: 37927234.
  16. Sjaastad AK, Jørgensen RB, Svendsen K. Exposure to polycyclic aromatic hydrocarbons (PAHs), mutagenic aldehydes and particulate matter during pan frying of beefsteak. *Occup Environ Med*. 2010 Apr; 67 (4): 228-32. doi: 10.1136/oem.2009.046144. Epub 2010 Feb 17. PMID: 20164502.
  17. Wei See S, Karthikeyan S, Balasubramanian R. Health risk assessment of occupational exposure to particulate-phase polycyclic aromatic hydrocarbons associated with Chinese, Malay and Indian cooking. *J Environ Monit*. 2006 Mar; 8 (3): 369-76. doi: 10.1039/b516173h. Epub 2006 Jan 19. PMID: 16528421.
  18. Sofuoglu SC, Toprak M, Inal F, Cimrin AH. Indoor air quality in a restaurant kitchen using margarine for deep-frying. *Environ Sci Pollut Res Int*. 2015 Oct; 22 (20): 15703-11. doi: 10.1007/s11356-015-4762-6. Epub 2015 May 29. PMID: 26022397.
  19. Lu F, Shen B, Yuan P, Li S, Sun Y, Mei X. The emission of PM2.5 in respiratory zone from Chinese family cooking and its health effect. *Sci Total Environ*. 2019 Mar 1; 654: 671-677. doi: 10.1016/j.scitotenv.2018.10.397. Epub 2018 Oct 30. PMID: 30448657.
  20. Lee YY, Park H, Seo Y, Yun J, et al. Emission characteristics of particulate matter, odors, and volatile organic compounds from the grilling of pork. *Environ Res*. 2020 Apr; 183: 109162. doi: 10.1016/j.envres.2020.109162. Epub 2020 Jan 22. PMID: 32018206.
  21. Sharma R, Balasubramanian R. Evaluation of the effectiveness of a portable air cleaner in mitigating indoor human exposure to cooking-derived airborne particles. *Environ Res*. 2020 Apr; 183: 109192. doi: 10.1016/j.envres.2020.109192. Epub 2020 Jan 27. PMID: 32062480.
  22. Kong HK, Yoon DK, Lee HW, Lee CM. Evaluation of particulate matter concentrations according to cooking activity in a residential environment. *Environ Sci Pollut Res Int*. 2021 Jan; 28 (2): 2443-2456. doi: 10.1007/s11356-020-10670-x. Epub 2020 Sep 4. PMID: 32888146.
  23. Lee IJ, Lee SG, Choi BH, Seo HK, Choi JH. Hazard Levels of Cooking Fumes in Republic of Korea Schools. *Saf Health Work*. 2022 Jun; 13 (2): 227-234. doi: 10.1016/j.shaw.2021.12.702. Epub 2022 Jan 3. PMID: 35664910; PMCID: PMC9142743.
  24. Padhi A, Bansal M, Habib G, Samiksha S, Raman RS. Physical, chemical and optical properties of PM2.5 and gaseous emissions from cooking with biomass fuel in the Indo-Gangetic Plain. *Sci Total Environ*. 2022 Oct 1; 841: 156730. doi: 10.1016/j.scitotenv.2022.156730. Epub 2022 Jun 14. PMID: 35714742.
  25. Lyu J, Shi Y, Chen C, Zhang X, Chu W, Lian Z. Characteristics of PM2.5 emissions from six types of commercial cooking in Chinese cities and their health effects. *Environ Pollut*. 2022 Nov 15; 313: 120180. doi: 10.1016/j.envpol.2022.120180. Epub 2022 Sep 16. PMID: 36122656.